


















COVID-19 vaccination schedule

Individuals not at greater risk of
serious disease from COVID-19 infection

Age (years)	0-11	12-15	16-17	18-39	40+
Primary course 	Vaccination not currently recommended	2 doses Pfizer BioNTech (Comirnaty®) >12 week interval ⁴		2 doses Pfizer (Comirnaty®) or Moderna (Spikevax®) ² 8 week interval	2 doses Pfizer (Comirnaty®) Moderna (Spikevax®) ² AstraZeneca (Vaxevria®) ⁵ 8 week interval
Booster 		 No Booster not recommended		 Yes Pfizer BioNTech (Comirnaty®) Moderna (Spikevax®) ² At least 3 month interval	
Infection 		Wait at least 12 weeks between COVID-19 infection and COVID-19 vaccination ⁴		Wait 4 weeks between COVID-19 infection and COVID-19 vaccination	

Individuals who are at greater risk of
serious disease from COVID-19 infection³

Age (years)	0-11	12-15	16-17	18-39	40+
<div>Primary course</div> <div></div>	<div>Vaccination not currently recommended</div>	<div>2 doses¹</div> <div>Pfizer BioNTech (Comirnaty®)</div> <div>8 week interval</div>		<div>2 doses¹</div> <div>Pfizer (Comirnaty®) or Moderna (Spikevax®)²</div> <div>8 week interval</div>	<div>2 doses¹</div> <div>Pfizer (Comirnaty®)</div> <div>Moderna (Spikevax®)²</div> <div>AstraZeneca (Vaxevria®)⁵</div> <div>8 week interval</div>
<div>Booster</div> <div></div>	<div></div>	<div> No</div> <div>Booster not recommended</div>	<div> Yes</div> <div>Pfizer BioNTech (Comirnaty®)</div> <div>At least 3 month interval</div>	<div> Yes</div> <div>Pfizer BioNTech (Comirnaty®) or Moderna (Spikevax®)²</div> <div>At least 3 month interval</div>	
<div>Infection</div> <div></div>	<div></div>	<div>Wait 4 weeks between COVID-19 infection and COVID-19 vaccination</div>			

¹3rd dose is advised if the person was severely immunocompromised at the time of first or second dose (refer to Green Book for details)

² The priming dose of Moderna Spikevax® COVID-19 Vaccine is 100 micrograms (0.5mL), The booster dose is 50 micrograms (0.25mL)

³ Health care, social care or care home staff, clinical risk groups, and those are severely immunosuppressed or their household contacts

⁴ This interval may be reduced to 8 weeks in periods of high incidence or where there was concern about vaccine effectiveness

⁵ Vaxevria should only be used in exceptional circumstances

